

Use this poster in conjunction with Quality Assurance Medication Label - Read and Match the Parts situation/task statement and the Medication Insert Parts Identification tags

Medication Insert

OMNIBIOTIC

(Hydrocillin in Aqueous Suspension)

For use in Beef Cattle, Lactating and Non-Lactating Dairy Cattle, Swine and Sheep

Read Entire Brochure Carefully Before Using This Product

For Intramuscular Use Only

Active Ingredients: Omnibiotic is an effective antimicrobial preparation containing hydrocillin hydrochloride. Each ml of this suspension contains 200,000 units of hydrocillin hydrochloride in an aqueous base.

Indications: **Cattle** - bronchitis, foot rot, leptospirosis, mastitis, metritis, pneumonia, wound infections. **Swine** - erysipelas, pneumonia.

Sheep - foot rot, pneumonia, mastitis; and other infections in these species caused by or associated with hydrocillin-susceptible organisms.

Recommended Daily Dosage

The usual dose is 2 ml per 100 lb of body weight given once daily. Maximum dose is 15 ml/day.

Body Weight	Dosage
100 lb	2 ml
300 lb	6 ml
500 lb	10 ml
750 lb or more	15 ml

Continue treatment for 1 to 2 days after symptoms disappear.

Caution: 1. Omnibiotic should be injected deep within the fleshy muscle of the neck or thigh. Do not inject this material in the hip or rump, subcutaneously, into a blood vessel, or near a major nerve because it may cause tissue damage. 2. If improvement does not occur within 48 hours, the diagnosis should be reconsidered and appropriate treatment initiated. 3. Treated animals should be closely observed for at least 30 minutes. Should a reaction occur, discontinue treatment and immediately administer epinephrine and antihistamines. 4. Omnibiotic must be stored between 2° and 8° C (36° to 46° F). Warm to room temperature and shake well before using. Keep refrigerated when not in use.

Warning: Milk that has been taken from animals during treatment and for 48 hours (4 milkings) after the last treatment must not be used for food. The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food.

How Supplied: Omnibiotic is available in vials of 100 ml.



Quality Assurance and Animal Care: Youth Education Program
This material is based upon work supported by Extension Service, United States Department of Agriculture, under special project number 93-EFSQ-4096.